



Sun Radio Interferometer Space Experiment (SunRISE)

Joseph Lazio, Justin Kasper, SunRISE Team

Solar Radio Bursts

Type II and III






SunRISE telescope

Solar Radio Bursts

Type II and III




frequency as coronal mass ejections expand into heliosphere Type II Radio Burst B Type III
Radio Burst 2000/02/27 07:42

Slowly descending in

Solar Radio Bursts

Type II and III



Slowly descending in frequency as coronal mass ejections expand into heliosphere


A Type II Radio Burst

B Type III Radio Burst


Rapidly drop in frequency as electron beams escape from active regions along open field lines

Science Implementation Concept

Synthetic Aperture!



Science Implementation



- First interferometer in space, first decametric-hectometric (DH) imaging
- Loose formation of six 6U form factor CubeSats in ~ 10 km sphere
- GEO+ Orbit (25 hr orbit period)
- DH radio receiver (0.1 MHz 20 MHz) with crossed 5 m dipole antennas
- Data collection in 0.655 ms bursts every 100 ms, select from 4096 frequencies
- Relative position knowledge to within 3 m and timing to ns
- Direct delivery to GEO+ as secondary payload
- 3 GB of data per week, all downloaded no bursts or triggers
- 2022 March Launch Readiness, six month Phase F

Space-based Radio Interferometer




Lift Long Wavelength Array-Owens Valley (or LWA1 or LOFAR or RAPID or ...) into space.

SunRISE telescope

Space-based Radio Interferometer

How hard can it be? • Power • Radiation environment • Reliability • Position and time knowledge • Data movement • Collisions(!)

Space-based Radio Interferometer



SunRISE telescope

Data Movement

Science Implementation Challenge

Observing at 10 MHz

Each spacecraft generates 160 Mbps, right?

- 10 MHz Nyquist sampled = 20 Msamples/s
- 2 polarizations
- 4 bit sampling

Commercial wireless routers advertise 1000+ Mbps

∴ No problem!

Data Movement

Science Implementation Challenge

Observing at 10 MHz

Each spacecraft generates 160 Mbps, right?

- 10 MHz Nyquist sampled = 20 Msamples/s
- 2 polarizations
- 4 bit sampling

Commercial wireless routers advertise 1000+ Mbps

∴ No problem!

Problem: spectrum allocations limit SunRISE to 2

Mbps

Solution: Reduce total bandwidth

Solution: Use DSN multi-spacecraft per aperture

Solution: Correlation done on ground

Correlation

Science Implementation Challenge

Position and time knowledge of spacecraft required

Sub-meter knowledge

Problem: 3 radios? Science, telecommunications, ranging


Mass, power, ...

Problem: Real-time orbit determination?

Solution: Use FX Correlator architecture coupled with GPS timing, dump spectra to ground, subsequent ground-based precise orbit determination and correlation

Collisions

Science Implementation Challenge



Collisions

Science Implementation Challenge


Spacecraft are 0.3 m in size (5 m antennas)

Separations >~ 1 km

∴ No problem!

Problem: (1 mm/s) × 1 week ~ 0.5 km

Solution: Orbital maintenance required



Space-based Low Radio Frequency Arrays

Astron. Astrophys. 195, 372-379 (1988)



A low frequency radio array for space

K.W. Weiler ¹, B.K. Dennison ^{1,2}, K.J. Johnston ¹, R.S. Simon ¹, W.C. Erickson ³, M.L. Kaiser ⁴, H.V. Cane ⁴, M.D. Desch ⁴, and L.M. Hammarstrom ¹

- ¹ E.O. Hulburt Center for Space Research, Naval Research Laboratory, Washington, DC 20375-5000, USA
- ² Virginia Polytechnic Institute and State University, Department of Physics, Blacksburg, VA 24061, USA
- 3 University of Maryland, Astronomy Program, College Park, MD 20742, USA
- 4 NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

Received August 4, accepted October 19, 1987

Summary. At the lowest radio frequencies (<30 MHz), the Earth's ionosphere transmits poorly or not at all. This relatively unexplored region of the electromagnetic spectrum is thus an area where high resolution, high sensitivity observations from space can open a new window for astronomical investigations. An array of free flying spacecraft which work as a coherent interferometer will be able to probe this frequency range. Operating from ~1 to ~30 MHz, such a telescope will extend astronomy from just above the ionospheric cutoff, where ground based observations can still be done, down to the fundamental physical limit where observations at still lower frequencies from within the Milky Way are impossible due to absorption by diffuse, ionized interstellar hydrogen.


telescopes. Only a few dedicated workers have continued to study the dekameter-hectometer wavelength radiation.

The most extensive investigations at the very lowest frequencies have been carried out with the Radio Astronomy Explorer (RAE) satellites 1 and 2 (Weber, Alexander, and Stone, 1971; Alexander and Novaco, 1974) in Earth and lunar orbit, respectively. They were launched at different dates and used as single survey antennas with their travelling wave V-antennas yielding only steradian resolution. Ground-based observations are normally confined to frequencies > 10 MHz during solar minimum or > 20 MHz during solar maximum, and only under special conditions at preferred locations does the ionosphere transmit radiation at frequencies as low as 2 to 5 MHz (Reber, 1968; Ellis

The scientific rewards of such a space mission are likely to be great. ... a low frequency telescope in space can ... study the impulsive low frequency emission from Jupiter and the Sun

The Astronomical Low Frequency Array

Viewing the Sun and Universe in a New Light



Dr. Dayton L. Jones Principal Investigator Mail code 238-332 Jet Propulsion Laboratory 4800 Oak Grove Drive Pasadena, CA 91109-0899 Phone: (818) 354-7774

Fax: (818) 393-6890 E-mail: dj@bllac.jpl.nasa.gov



Dr. Charles Elachi Space and Earth Sciences Directorate Mail code 180-704 Jet Propulsion Laboratory

4800 Oak Grove Drive Pasadena, CA 91109-0899 Phone: (818) 354-5673 Fax: (818) 354-2946

E-mail: Charles.Elachi@jpl.nasa.gov

RESTRICTION ON USE AND DISCLOSURE OF PROPOSAL AND QUOTATION INFORMATION

The information contained in all pages of this proposal or quotation constitutes a trade secret and/or information that is commercial or financial and confidential or privileged. It is furnished to the Government in confidence with the understanding that it will not, without permission of the offeror, be used or disclosed for other than evaluation purposes; provided, however, that in the event a contract is awarded on the basis of this proposal or quotation, the Government shall have the right to use and disclose this information to the extent provided in the contract. This restriction does not limit the Government's right to use of access this information if obtained from another over without restriction.

Sun Radio Interferometer Space Experiment

Space-based Radio Interferometer

Dawn of space-based arrays Small spacecraft enabling technology SunRISE in Phase A

Exciting science


Fundamental plasma physics in the heliosphere, planetary magnetospheres, nearby pulsars, ...

Scaling to much larger arrays may yet require technology development

Miniaturization, new data movement approaches, robust real-time position and timing knowledge, on-orbit correlation

backup

Concept of Operations



Solar and Space Physics


Goal 1. Determine the origins of the Sun's activity and predict the variations in the space environment.

Goal 2. Determine the dynamics and coupling of Earth's magnetosphere, ionosphere, and atmosphere

Goal 3. Determine the interaction of the Sun with the solar system and the interstellar medium.


Goal 4. Discover and characterize fundamental processes that occur both within the heliosphere and throughout the universe.

SHP Panel Goal 3. Determine how magnetic energy is stored and explosively released.



SunRISE Objective #1

The Acceleration of Solar Energetic Particles




Discriminate competing hypotheses for the generation of solar energetic particles by measuring the location and morphology of associated Type II radio emissions.

Major solar energetic particle events proceeded (95% of time) by Type II radio burst below 15 MHz


SunRISE Objective #2

The Release of Solar Energetic Particles into Space



Discriminate competing hypotheses for the variable magnetic connection between active regions and the inner heliosphere by reconstructing magnetic field lines associated with Type III radio bursts.

CME Evolution and Particle Acceleration



Ground-based measurements limited to r ~ 2 R_☉

Limited frequency range cannot track evolution, limits extent to which radio-optical images can be aligned /correlated

CME (white light)

80 MHz Gauribidanur image

SDO-AIA 193 Å and SOHO-LASCO C2 images

Science Implementation

6 6U spacecraft

Observing 0.1 MHz—25 MHz

Located in supersynchronous GEO orbit (a.k.a. GEO graveyard)

Observing solar Type II and Type III bursts

NASA/Heliophysics Announcement of Opportunity	c. 2016 July 20
SunRISE proposal submitted (NASA/Heliophysics SALMON-2 PEA Q/MOO SCM)	2016 October 14
SunRISE selected	2017 July 28
SunRISE Concept Study Report due	2018 July 30
Launch	c. 2022 July


Breaking News!

July 28, 2017

NASA Selects Proposals to Study Sun, Space Environment








DHFR Launch August 25!




SunRISE

Schedule



Sun Radio Imaging Space Experiment

Mission Concept



Position Determination

